



This label contains the most current product information and may differ from the information on the product package.

Vazalore™ 325 mg Capsules

Drug Facts	
Active ingredient (in each capsule)	Purposes
Aspirin (NSAID)* 325 mg *nonsteroidal anti-inflammatory drug	Pain reliever/fever reducer
Uses	
<ul style="list-style-type: none"> ■ for the temporary relief of minor aches and pains associated with: <ul style="list-style-type: none"> ■ headache ■ backache ■ muscular aches ■ a cold ■ toothache ■ minor pain of arthritis ■ premenstrual and menstrual cramps ■ temporarily reduces fever ■ ask your doctor about other uses for aspirin capsules 	
Warnings	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) <p>This product contains soy.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed 	
<p>Do not use if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.</p>	
Ask a doctor before use if	
<ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma 	
Ask a doctor or pharmacist before use if you are	
<ul style="list-style-type: none"> ■ taking a prescription drug for diabetes, gout or arthritis 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts for more than 3 days ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms occur. These could be signs of a serious condition. 	
<p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	
Directions	
<ul style="list-style-type: none"> ■ drink a full glass of water with each dose ■ adults and children 12 years and over: <ul style="list-style-type: none"> ■ take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist ■ do not exceed 12 capsules in 24 hours ■ children under 12 years: consult a doctor 	
Other information	
<ul style="list-style-type: none"> ■ save carton for full directions and warnings ■ store at 15-30°C (59-86°F) ■ avoid excessive heat above 40°C (104°F) ■ do not use if the blue band around the capsule is missing or broken 	
Inactive ingredients anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, lecithin, oleic acid, pharmaceutical ink, potassium chloride, soybean oil, titanium dioxide	
Questions or comments?	
Call 1-888-842-3567 9:00 AM - 7:00 PM ET, Monday - Friday	

Revision 07-27-21