



This label contains the most current product information  
and may differ from the information on the product package.

## Vazalore™ 325 mg Capsules

### Drug Facts

#### Active ingredient (in each capsule)

Aspirin (NSAID)\* 325 mg ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Purposes

#### Uses

- for the temporary relief of minor aches and pains associated with:
  - headache    ■ backache    ■ muscular aches    ■ a cold    ■ toothache
  - minor pain of arthritis    ■ premenstrual and menstrual cramps
- temporarily reduces fever
- ask your doctor about other uses for aspirin capsules

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives    ■ facial swelling    ■ shock    ■ asthma (wheezing)

This product contains soy.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer.

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

#### Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

#### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint    ■ vomit blood    ■ have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts for more than 3 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms occur. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- drink a full glass of water with each dose
- adults and children 12 years and over:
  - take 1 or 2 capsules every 4 hours or 3 capsules every 6 hours while symptoms persist
  - do not exceed 12 capsules in 24 hours
- children under 12 years: consult a doctor

#### Other information

- save carton for full directions and warnings
- store at 15-30°C (59-86°F)
- avoid excessive heat above 40°C (104°F)
- do not use if the blue band around the capsule is missing or broken

**Inactive ingredients** anhydrous citric acid, carrageenan, colloidal silicon dioxide, FD&C Blue #1, hypromellose, lecithin, oleic acid, pharmaceutical ink, potassium chloride, soybean oil, titanium dioxide

#### Questions or comments?

Call 1-888-842-3567 9:00 AM - 7:00 PM ET, Monday - Friday

Revision 07-27-21